# Enablers and barriers to workplace breastfeeding in the Armed Forces: a systematic review

Hannah Taylor 💿

#### ► Additional material is published online only. To view, please visit the journal online (http://dx.doi.org/10.1136/ bmjmilitary-2020-001724).

#### Correspondence to

Dr Hannah Taylor, Army Medical Services, Camberley, UK; hannah.taylor43@nhs.net

Received 25 November 2020 Revised 12 January 2021 Accepted 13 January 2021 Published Online First 16 February 2021 **ABSTRACT Introduction** The UK has no legislation protecting employees' access to breastfeeding facilities. Without specific breastfeeding policy, provisions to access workplace facilities can be inconsistent and negatively impact employees' breastfeeding duration, retention and morale, particularly servicewomen who work in varied and demanding military environments. This is an important policy area for the British Army to retain talented and trained soldiers.

**Methods** Using Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement principles, PubMed, Embase, CINAHL and Pro-Quest Databases were searched for studies relevant to accessing appropriate breastfeeding facilities in UK workplaces and highincome countries' Armed Forces. Factors acting as barriers and enablers to accessing facilities were identified. UK government and Armed Forces' websites were searched for grey literature on existing policies and guidance for accessing facilities.

**Results** Barriers and enablers to access from 16 studies were described by three thematic areas: attitudes to breastfeeding, facility provisions and use of facilities. Factors which employers could influence included specific breastfeeding policy, universal workplace education, existence of suitable facilities and individualised breastfeeding plans. The key areas for policy development identified were clearly defined responsibilities; individualised risk assessments and breastfeeding plans; appropriate, but flexible, facility provision and access; signposting of relevant workplace accommodations; and physical fitness provisions.

**Conclusions** Five recommendations are presented: development, implementation and evaluation of breast-feeding policy; universal workplace breastfeeding education; the need for breastfeeding risk assessments and plans based on individual breastfeeding practice; written minimal and ideal standards for breastfeeding facilities and access, which considers workplace locations; and exceptions from deployment and physical fitness testing.

## **INTRODUCTION**

The World Health Organisation (WHO) recommends exclusive breastfeeding for six months and continuation, alongside solids, to the age of two years.<sup>1 2</sup> Breastfeeding benefits infant<sup>3 4</sup> and maternal long-term health,<sup>4-6</sup> employee presenteeism<sup>7</sup> and retention.<sup>8 9</sup> However, for many complex demographic,<sup>10</sup> cultural<sup>11</sup> and socioeconomic reasons,<sup>12 13</sup> United Kingdom (UK) breastfeeding rates are among the world's lowest.<sup>14</sup> The UK is one of 15/56 high-income countries (HICs) without legislation guaranteeing suitable workplace

## Key messages

- ⇒ Facilitating breastfeeding on return to work (RTW) benefits the maternal and infant physical and mental health, and can improve employee presenteeism, loyalty and retention.
- ⇒ More research and evaluation of breastfeeding servicewomen in the United Kingdom (UK) Armed Forces or other militaries around the world are needed to better support this population.
- ⇒ A single, easily accessible breastfeeding-specific policy or guidance document that clearly states minimum and optimal standards is required for both servicewomen and commanders.
- ⇒ Facility access and appropriateness are improved by ensuring individual risk assessments and breastfeeding plans, which account for individual circumstances and career group requirements, are undertaken.
- ⇒ Organisational cultural acceptability of breastfeeding is influenced by universal workplace, commander and medical chain education covering breastfeeding on RTW specific to the setting.
- ⇒ Effective peer support and empowerment of breastfeeding servicewomen to address the unique challenges faced could be achieved through a military-specific breastfeeding network.

facilities or breastfeeding breaks.<sup>15</sup> <sup>16</sup> The only relevant legislation states: '*Suitable facilities shall be provided* (for) *a nursing mother to rest.*<sup>17</sup> This must not be the toilets.<sup>18</sup> <sup>19</sup> Consequently, access to suitable workplace breastfeeding facilities varies significantly. These uncertainties can delay maternal return to work (RTW) or shorten breastfeeding duration.<sup>12</sup> <sup>20</sup> Societal expectations of service-women, periods of separation, physical fitness requirements, deployments and hazardous environmental exposures<sup>21</sup> make Armed Forces RTW breastfeeding even more complex.

All nations' servicewomen have similar roles, but differences in working conditions and breastfeeding provisions mean recommendations<sup>22</sup> from other militaries are not always transferable to the British Army. Although UK triservice policy advocates private safe spaces with breastmilk storage,<sup>23</sup> in reality, breastfeeding expectations and provisions depend on individual commanders' approaches. Such uncertainties may contribute to decisions to terminate service or resentfully cease

Check for updates

© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Taylor H. *BMJ Mil Health* 2023;**169**:373–384.

Table 1	Study characteristics						
	Author	Publication year	Study design	Studies reviewed	Workplace	Relevant main outcomes	Critical appraisa score
HICs' Armed Forces	Croft <sup>22</sup>	1995	Literature review	16	British Army	Occupational environmental exposures health risks to breastfeeding mothers and their infants	0.7
				Sample size	Population		
	Bales et al <sup>35</sup>	2012	Cross-sectional telephone survey	43 servicewomen 211 service spouses	USAF base mothers	<ul> <li>Barriers and facilitators to meeting personal breastfeeding goals.</li> </ul>	0.7
	Bell and Ritchie <sup>36</sup>	2003	Cross-sectional qualitative	85	US military TRICARE facility stakeholders	<ul> <li>Impact of workplace lactation support services on breastfeeding duration.</li> </ul>	0.6
	Bristow <sup>43</sup>	1999	Cross-sectional qualitative	4	US servicewomen, all branches	<ul> <li>Workplace breastfeeding facilitators and barriers.</li> </ul>	0.8
	Harlow <sup>44</sup>	1998	Cross-sectional written survey	64	US servicewomen, all branches	<ul> <li>Facilitators and barriers to breastfeeding initiation and workplace continuation.</li> </ul>	0.9
	Martin <i>et al<sup>37</sup></i>	2015	Cross-sectional online survey	318	US servicewomen, all branches	<ul> <li>Perceptions of breastfeeding in the workplace.</li> <li>Workplace Breastfeeding Support Scale score.</li> </ul>	0.8
	Sleutel <sup>38</sup>	2012	Cross-sectional qualitative	1	US Army servicewomen	<ul> <li>Breastfeeding experiences in firm-based and deployed locations.</li> </ul>	0.4
	Stevens and Janke <sup>39</sup>	2003	Cross-sectional qualitative	9	USAF servicewomen	<ul> <li>Workplace breastfeeding experiences.</li> </ul>	0.7
	Stewart <sup>40</sup>	2015	Cross-sectional online survey	152	Australian Defence Force (ADF) servicewomen	<ul> <li>Impact of differing RTW and breastfeeding models on breastfeeding duration.</li> <li>Causes of RTW breastfeeding cessation.</li> </ul>	0.9
	Uriell <i>et al</i> <sup>41</sup>	2009	Cross-sectional online survey	7121 personnel 2195 breastfeeding servicewomen	US Naval personnel and breastfeeding servicewomen	<ul> <li>Causes of RTW breastfeeding cessation.</li> </ul>	0.9

BMJ Mil Health: first published as 10.1136/bmjmilitary-2020-001724 on 16 February 2021. Downloaded from http://militaryhealth.bmj.com/ on April 27, 2024 by guest. Protected by copyright.

Continued

Table 1 Continued

	Author	Publication year	Study design	Studies reviewed	Workplace	Relevant main outcomes	Critical appraisal score
UK	Gatrel <sup>30</sup>	2007	Cross-sectional qualitative	20	Professional or managerial breastfeeding mothers	<ul> <li>Workplace breastfeeding experiences.</li> <li>Impact of personal perceptions of working environment on breastfeeding behaviours.</li> </ul>	0.8
	Hawkins <i>et al</i> <sup>31</sup>	2007	Cohort study	6917	Employed and unemployed mothers	<ul> <li>Breastfeeding duration by work pattern.</li> </ul>	0.9
	Kosmala-Anderson and Wallace <sup>32</sup>	2006	Cross-sectional online survey	46	Postpartum public-sector employees	<ul> <li>Experiences of workplace breastfeeding support.</li> </ul>	0.6
	Skafida <sup>33</sup>	2011	Cohort study	5127	Employed and unemployed Scottish mothers	<ul> <li>Causes of cessation at 10 months.</li> </ul>	0.8
	Wallace <i>et al</i> <sup>34</sup>	2008	Cross-sectional online survey	296	Breastfeeding employees	<ul> <li>RTW breastfeeding experiences.</li> <li>Supportive factors and barriers to effective workplace breastfeeding.</li> </ul>	0.7
	Zilanawala <sup>42</sup>	2017	Cohort study	17 397	Employed and unemployed mothers receiving child benefit	<ul> <li>Breastfeeding rates at 9 months by work pattern.</li> </ul>	0.7

HIC, high-income country; RTW, return to work; USAF, US Airforce.

breastfeeding.<sup>24</sup> The proportion of trained and talented women, who the British Army need to retain,<sup>25</sup> is ever increasing, and with 6.4% taking maternity leave annually,<sup>26</sup> breastfeeding policy is required. This review identifies barriers and enablers to accessing appropriate workplace breastfeeding facilities and the relevance of these factors to the British Army.

## **METHODOLOGY**

A literature review was undertaken in 2017 in accordance with ethics approval. Standardised Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement<sup>27</sup> principles were adhered to. To account for occupational circumstances and UK workplace legislation, the search strategy combined UK workplaces and English-speaking HIC military breastfeeding studies, existing policy and guidance. Databases searched included the Cochrane Library, CINAHL, PubMed, Embase and PRO-Quest Military Collection (Supplement S3). The World Bank's definition of an HIC, as one with a per capita income of \$12 476 or more, was used. The UK Government and Britishspeaking Defence sites were searched for publicly available grey literature (Supplement S4). To improve search sensitivity and to ensure comprehensive study inclusion, common synonyms and Boolean operators for key concepts were used; Breastfeeding, Employment, Barrier, Enabler, Facilities and Military, Search specificity was improved following a pilot, by making 'breast' a mandatory term (Supplements S1 & S2).

Inclusion criteria for UK workplace studies were English language and peer-reviewed journal publication post-2006. HICs Armed Forces studies were limited; therefore, all publicly available studies and theses, of any age, with a formal review and ethics process, were included. For English language policies and guidance, newest versions were sought. Opinion pieces, news stories, informal advice, social media recommendations, and studies focusing on healthcare professionals (HCPs) or partner outcomes were excluded. All search results details were exported into Endnote V.X7.5 and duplicates were removed. Remaining abstracts and documents were scanned for suitability, and results are displayed in Tables 1–4.

Study quality was assessed using validated Critical Appraisal Skills Programme tools<sup>28</sup> for literature reviews, cohort and qualitative studies; the Strengthening the Reporting of Observational Studies in Epidemiology checklist assessed bias in cross-sectional studies (Supplement S5).<sup>29</sup> As each tool assessed a differing number of elements, the mean average score was calculated; each element scored fully (1.0), partially (0.5) or not (0) achieved. Mean scores were used to grade study quality as unsatisfactory (0–0.4), satisfactory (0.41–0.7) or good (0.71–1.0).

## RESULTS

Figure 1 illustrates primary search strategy results.<sup>27</sup> Table 1 details the 16 included studies' characteristics, risk of bias and quality. Due to limited study availability, one unsatisfactory

			Croft <sup>22</sup>	Bales et al <sup>35</sup>	Bell nd Ritchie <sup>36</sup>	Bristow <sup>43</sup>	Harlow <sup>44</sup>	Martin et al <sup>37</sup>	Sleutel <sup>38</sup>	Stevens and Janke <sup>39</sup>	Stewart <sup>40</sup>	Uriell et al <sup>41</sup>	Gatrel <sup>30</sup>	Hawkins et al <sup>31</sup>	Kosmala-Anderson and Wallave <sup>32</sup>	Skafida <sup>33</sup>	Wallace et al <sup>34</sup>	Zilanawala <sup>42</sup>	Total
ides to	Initiation and	Maternal attitudes		×		×	×	×	×	×			×	×	×	×		×	=
tfeeding	duration	Antenatal knowledge about workplace breastfeeding			×		×								×		×		4
	Acceptability	Knowledge on breastfeeding benefits mother, infant and employer			×		×	×	×					×	×		×		2
		Protective breastfeeding policies			×	×		×	×					×	×		×		7
		Workplace breastfeeding support			×	×	×	×		×			×	×	×		×	×	10
	Affordability	Perceived long-term benefits for employer			×	×		×	×										4
		Ability to make informed choices about career impact	×														×		2
ion of priate es	Expectations	Universal education on managing workplace breastfeeding		×	×	×		×	×				×	×	×		×		6
		Military-specific breastfeeding class		×	×		×												m
		Individualised breastfeeding plan		×		×		×	×	×							×	×	7
	Availability	Accommodations to access facilities external to workplace				×		×			×				×	×	×	×	٢
		Legislation or policy requirement			×										×				2
		Workplace facility exists			×	×									×		×		4
		On-site nursery													×		×		2
	Suitability	Private safe space			×	×				×		×	×		×		×	×	∞
		Suitable amenities			×	×			×			×			×		×		9
		Refrigerated storage and logistical support							×						×		×	×	4
		Provision of enabling equipment			×	×		×	×	×							×		9

BN
2
_ii ⊤
leal
5: f
irst
pub
lish
ed a
as 1
0.1
136,
/bm
mili
tary
-20
20-0
0017
724
DN,
16 F
ebr
uary
/ 20
21.
Dov
vnlo
ade
ăfr
m
http
://m
litar
yhe
alth
.bm
j.co
m c
n A
pril
27,
202
4 by
ng /
est.
Pro
vtect
ted l
by c
ору
righ

Systematic review

	Total	m	m	7	2	4	4	9	m	7	9
	ilanawala <sup>42</sup>										
	allace al <sup>34</sup> Z			×			×	×	×	×	
	la <sup>33</sup> et			×				×		×	
	ר Skafic							×			
	Kosmala-Anderso and Wallave <sup>32</sup>	×			×	×		×		×	
	Hawkins et al <sup>31</sup>										
	Gatrel <sup>30</sup>									×	
	Uriell <i>et al</i> <sup>41</sup>			×	×	×				×	
	Stewart <sup>40</sup>			×							
	Stevens and Janke <sup>39</sup>			×			×	×			
	Sleutel <sup>38</sup>	×	×				×		×		
	Martin et al <sup>37</sup>			×			×	×	×	×	×
	Harlow <sup>44</sup>										×
	Bristow <sup>43</sup>			×		×				×	×
	3ell nd Ritchie <sup>36</sup>										
	Bales   et al <sup>35</sup>	î	×								×
	Croft <sup>22</sup>		×								
		Timely knowledge of facilitates	Actioned risk assessment and breastfeeding plan	Role allowing work schedule self- management	Unrestricted access to facilities	Formally agreed breastfeeding breaks	Advance planning for changes to working location or pattern	Flexible-working provisions	Workplace breastfeeding support network	Workplace support to access facilities	Medical chain support for breastfeeding
Continued		Knowledge		Access					Support		
Table 2 (		Use of facilities									

 .

Syster	natic	revie	w	

Antiolity         Antiolity <t< th=""><th>Ba</th><th>rriers to ac</th><th>ccessing appropria</th><th>ate facili</th><th>ties</th><th></th><th></th><th></th><th></th><th></th><th>Stevens</th><th></th><th></th><th></th><th></th><th></th><th></th><th></th><th></th><th></th><th></th></t<>	Ba	rriers to ac	ccessing appropria	ate facili	ties						Stevens										
411 <t< th=""><th></th><th></th><th></th><th>Croft<sup>22</sup></th><th>Bales et al<sup>35</sup></th><th>Bell nd Ritchie<sup>36</sup></th><th>Bristow<sup>43</sup></th><th>Harlow<sup>44</sup></th><th>Martin et al<sup>37</sup></th><th>Sleutel<sup>38</sup></th><th>and Janke<sup>39</sup></th><th>Stewart<sup>40</sup></th><th>Uriell <i>et al</i><sup>41</sup> Gatı</th><th>rell<sup>30</sup> et à</th><th>wkins Kosn a/<sup>31</sup> and</th><th>ıala-Anderson Nallace<sup>32</sup></th><th>Skafida<sup>33</sup></th><th>Wallace et al<sup>34</sup></th><th>Zilanawala<sup>42</sup></th><th>Total</th><th></th></t<>				Croft <sup>22</sup>	Bales et al <sup>35</sup>	Bell nd Ritchie <sup>36</sup>	Bristow <sup>43</sup>	Harlow <sup>44</sup>	Martin et al <sup>37</sup>	Sleutel <sup>38</sup>	and Janke <sup>39</sup>	Stewart <sup>40</sup>	Uriell <i>et al</i> <sup>41</sup> Gatı	rell <sup>30</sup> et à	wkins Kosn a/ <sup>31</sup> and	ıala-Anderson Nallace <sup>32</sup>	Skafida <sup>33</sup>	Wallace et al <sup>34</sup>	Zilanawala <sup>42</sup>	Total	
Indefinition </td <td>Initiation</td> <td>and</td> <td>Maternal attitudes</td> <td></td> <td></td> <td></td> <td>×</td> <td>×</td> <td>×</td> <td>×</td> <td>×</td> <td>×</td> <td>×</td> <td>×</td> <td></td> <td></td> <td>×</td> <td></td> <td></td> <td>6</td> <td></td>	Initiation	and	Maternal attitudes				×	×	×	×	×	×	×	×			×			6	
Independencie       Independencie <thindependencie< th=""> <thindependencie< t<="" td=""><td>duration</td><td></td><td>Practical difficulties expressing</td><td></td><td>×</td><td></td><td>×</td><td>×</td><td></td><td></td><td>×</td><td></td><td>×</td><td></td><td>×</td><td></td><td>×</td><td></td><td></td><td>7</td><td></td></thindependencie<></thindependencie<>	duration		Practical difficulties expressing		×		×	×			×		×		×		×			7	
Ad foldigit       X <td< td=""><td>Accepta</td><td>oility</td><td>Negative perceptions of workplace breastfeeding</td><td></td><td></td><td>×</td><td>×</td><td></td><td></td><td></td><td>×</td><td>×</td><td>×</td><td></td><td>×</td><td></td><td></td><td></td><td></td><td>9</td><td></td></td<>	Accepta	oility	Negative perceptions of workplace breastfeeding			×	×				×	×	×		×					9	
igo downlote       x <t< td=""><td></td><td></td><td>Lack of knowledge about breastfeeding at work</td><td></td><td></td><td></td><td></td><td>×</td><td>×</td><td></td><td></td><td></td><td>×</td><td></td><td>×</td><td></td><td></td><td>×</td><td></td><td>ъ</td><td></td></t<>			Lack of knowledge about breastfeeding at work					×	×				×		×			×		ъ	
10       Inductor       x       x         Andread       X       X       X </td <td></td> <td></td> <td>Lack of workplace support</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>×</td> <td></td> <td>×</td> <td></td> <td>×</td> <td></td> <td></td> <td></td> <td>×</td> <td>×</td> <td></td> <td>ß</td> <td></td>			Lack of workplace support						×		×		×				×	×		ß	
Cutrantication         X         X         X           Reprint         X         X         X         X	Affordat	oility	Financial costs of facilities										×							-	
Periodiant       x       x       x       x         information       information       x       x       x       x         information       information       x       x       x       x         information       x       x       x       x       x       x         information       x       x       x       x       x       x       x         information       x       x       x       x       x       x       x       x         information       x       x       x       x       x       x       x       x         information       x       x       x       x       x       x       x       x       x         information       x </td <td></td> <td></td> <td>Career ramifications</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>×</td> <td></td> <td>×</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>m</td> <td></td>			Career ramifications								×		×							m	
100       Underformediate       x			Perceived impact on workplace effectiveness			×			×		×		×		×					2	
Outlong to the formation of the fo	Expecta	tions	Unrealistic expectations for provisions				×			×										2	
ID       Monstant facility       x       x       x       x         Attail or exercted priving location of purple sectors       x       x       x       x       x         Attail or exercted priving location of purple sectors       x       x       x       x       x       x         Attail or exercted purple sectors       x       x       x       x       x       x       x       x         Attail or exercted purple sectors       x </td <td></td> <td></td> <td>Lack of HCP workplace breastfeeding advice</td> <td></td> <td></td> <td></td> <td>×</td> <td>×</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>×</td> <td></td> <td></td> <td>×</td> <td></td> <td>4</td> <td></td>			Lack of HCP workplace breastfeeding advice				×	×							×			×		4	
Autor oppeted       X       X	Availabi	lity	Non-existent facility		×	×							×		×			×		5	
Image: A control decision       x<			Actual or expected changes to workload, working location or pattern		×						×								×	m	
Update       Let of policy of the solution of the soluticon of the solution of the solution of the sol			Lack of workplace breastfeeding research						×				×							2	
Watchelocation       X			Lack of policy or awareness of it			×				×			×		×			×		2	
Haradus x x x x x x x x x x x x x x x x x x x	Suitabili	A1	Unsuitable location for breastfeeding		×	×	×		×		×		×		×			×		œ	
Lack of suitable x x x x x x x x x x x x x x x x x x x			Hazardous occupational environmental exposures	×						×			×		×					4	
Lack of breastmilk x x x x x x transporting to the infant transporting tran			Lack of suitable amenities		×	×	×		×	×			×		×			×		œ	
			Lack of breastmilk storage or logistical challenges transporting it to infant							×	×		×		×					4	
																				Continu	pani

q									Stevens								
		Croft <sup>22</sup>	Bales et al <sup>35</sup>	Bell nd Ritchie <sup>36</sup>	Bristow <sup>43</sup>	Harlow <sup>44</sup>	Martin e <i>t al<sup>37</sup></i> Sl	a leutel <sup>38</sup> J	and anke <sup>39</sup> Si	tewart <sup>40</sup>	Uriell <i>et al<sup>41</sup> G</i> a	H: Itrell <sup>30</sup> et	awkins K t <i>al</i> ³1 a	osmala-Anderson nd Wallace <sup>32</sup>	N Skafida <sup>33</sup> e	allace <i>al</i> <sup>34</sup> Zilanawala <sup>42</sup>	Total
Lack o availat	f knowledge of ble facilities			×	×								×			×	4
Male work	-dominated place					^	×				×				×		m
Supe unav brea	ervisor ware mother is istfeeding			×													-
Acc	ess to facilities sically restricted						×	×	ž		×					×	4
ы Э	cupational stress or gh workload		×			~	×	×	ž		×				×	×	9
La	ck of time			×	×		×	×	ý		×		×				9
p R	on-standard working tterns							×	,		×				×	×	4
₽š	lerance of negative orkplace behaviours										×						2
Se ex ne	paration from ternal support twork		×		×	×		×	~		×	×				×	7
S C	ack of workplace upport				×	~	×	×	×		×		×		×	×	6

# Systematic review

quality study was included; however quality was considered in the discussion. Fourteen studies were peer-reviewed<sup>30-41</sup> (1995– 2017),<sup>22 42</sup> including two US theses (1998–1999).<sup>43 44</sup> Six studied UK mothers<sup>30-34 42</sup> and 10 HIC servicewomen,<sup>22 35-44</sup> of which 80% were US-centric. Differences in gender, ethnicity and role composition made studies less representative of a contemporaneous British Army, demonstrating why a review was required.

Study design varied significantly and included one literature review of otherwise not-included studies<sup>22</sup>; three large UK<sup>31 33 42</sup> population cohort studies; seven quantitative crosssectional studies, using online, <sup>32 34 37 38 41</sup> written<sup>44</sup> or telephone surveys<sup>35</sup>; and five cross-sectional qualitative studies, using interviews<sup>30 36 39</sup> servicewomen<sup>30 38 39 43</sup> or stakeholder<sup>36</sup> narratives.<sup>38 43</sup> Only three recognised limitations of recall bias in cross-sectional studies, especially where workplace recollections may be corrupted by overall breastfeeding experience. Only two addressed<sup>33 41</sup> selection bias and temporality<sup>8</sup> of including only currently working mothers<sup>30 32 34 35 37-41 43</sup>; excluded mothers may delay RTW to breastfeed longer. UK civilian employees may also differ from servicewomen; characteristics which drive enlistment may also influence responses to breastfeeding barriers.

Differing study methodologies, sizes (range 1–17 597 mothers) and outcomes measures limited comparability and accurate quantification of factor dominance or breastfeeding outcome. Additionally, no single concept of access model<sup>45</sup> could be used to explain all identified factors or their interconnectivity. However, inclusion of multiple complementary study designs offered scope to identify a more comprehensive range of enabling factors and barriers (Tables 2 and 3).<sup>46</sup> Findings were based only on factors which emerged from thematic analysis of the included studies and therefore may not be exhaustive. All themes interacted through feedback loops within a complex system. While some connections were explored, conclusions drawn may oversimplify, may not fully explain or may underestimate the root cause factors influencing access to breastfeeding facilities.

Table 4 details characteristics of the 16 RTW policy and guidance documents. Table 5 identifies the five key areas from policy analysis, in which recommendations can be made: employer, employee and medical responsibilities; individualised risk assessments and breastfeeding plans; requirements for breastfeeding facility provision and access; provisions for RTW physical fitness; and workplace accommodations.

## Attitudes

The most common enabler to accessing breastfeeding facilities, identified by six military<sup>35</sup> <sup>37–39</sup> <sup>43</sup> <sup>44</sup> and five UK studies,<sup>30–33</sup> <sup>42</sup> was maternal motivation to reach a predetermined goal. Maternal disclosure of breastfeeding status, as well as behaviour actively seeking access to breastfeeding facilities on RTW, was positively influenced by actual or perceived supervisor, military HCP<sup>35–38</sup> <sup>43</sup> <sup>44</sup> or colleague support.<sup>30</sup> <sup>32</sup> <sup>34</sup> <sup>36</sup> <sup>37</sup> <sup>39</sup> <sup>42–44</sup> Workforce acceptance was strongly influenced by knowledge of the benefits of breastfeeding to the employer.<sup>37</sup> <sup>38</sup> <sup>43</sup>

Lack of support, <sup>33 34 37 39 41</sup> workplace pressure to stop and separation from external breastfeeding support networks all reduced breastfeeding duration. <sup>31 35 39 41-44</sup> Hostile behaviours<sup>30-41</sup> were fuelled by limited employer education and subsequent misconceptions. <sup>30 32 34 37 41</sup> These included maternity leave being the only acceptable time and place to breastfeed, <sup>40 43</sup> support for breastfeeding conveying a subversive feminine organisational image<sup>32</sup> and negatively impacting workforce effectiveness. <sup>30 32 33 39 43</sup> Military studies identified concerns that breastfeeding could threaten operational deployability and capability, but all concluded

Owner	Document type and target audience	Publication year	Main purpose
HSE <sup>18</sup>	Guide for UK mothers during pregnancy or on RTW	2013	Informing employees of health and safety at work legislation protecting pregnant and breastfeeding mothers.
ACAS <sup>19</sup>	Guide for UK employees and employers on accommodating workplace breastfeeding	2017	<ul> <li>Guidance on managing workplace breastfeeding requests and the legislation and good practice recommendations for facilities</li> </ul>
NHS <sup>51</sup>	Information for breastfeeding mothers on RTW	2008	<ul> <li>Information for mothers about continuing breastfeeding on RTW</li> <li>A resource to show employers.</li> </ul>
MOD <sup>59</sup>	Leave policies for UK service personnel	2016	<ul> <li>Policy on all leave types, including maternity leave.</li> </ul>
MOD 43	Health and safety in defence policy	2016	<ul> <li>Policy on health and safety requirements for service personnel</li> </ul>
The Royal Navy <sup>60</sup>	Personnel management policies	2016	<ul> <li>Policy on personnel management, including pregnant and postpartum servicewomen.</li> </ul>
RAF 52	Guide for pregnant and postpartum servicewomen	2014	<ul> <li>Handbook for pregnant and postpartum servicewomen on RTW.</li> </ul>
RAF 53	Pregnancy and postpartum guide for CoC	2017	Handbook for pregnant and postpartum servicewomen's CoC.
USAF <sup>54</sup>	Medical care management instruction	2017	<ul> <li>Guidance on the medical care and management, including of pregnant, postpartum and breastfeeding servicewomen.</li> </ul>
USAF <sup>44</sup>	COC memorandum for deployment and physical exercise deferment	2015	<ul> <li>Guidance on extending posting, deployment or physical assessment deferrals.</li> </ul>
US DOD <sup>48</sup>	US Army memorandum to breastfeeding support policy for servicewomen and the CoC	2016	<ul> <li>Policy on provisions for access to appropriate breastfeeding facilities.</li> <li>Outlines individual, CoC and medical chain responsibilities.</li> <li>Provides paperwork for an individualised breastfeeding plan.</li> </ul>
DOD <sup>42</sup>	US Army breastfeeding support plan for servicewomen and the CoC	2016	<ul> <li>Universal educational on breastfeeding.</li> <li>Guidance for supporting workplace breastfeeding.</li> <li>Provides paperwork for a workplace breastfeeding plan.</li> </ul>
US Coast Guard <sup>55</sup>	Pregnancy and postpartum policy	2016	<ul> <li>Policy on managing pregnant and postpartum servicewomen.</li> </ul>
DOD 56	US Marines pregnancy and postpartum policy	2004	<ul> <li>Policy on managing pregnant and postpartum servicewomen.</li> </ul>
DOD 57	US Naval pregnancy and postpartum instruction	2007	<ul> <li>Policy on managing pregnant and postpartum servicewomen.</li> </ul>
RAAF <sup>49</sup>	Line managers handbook for managing breastfeeding mothers	2014	<ul> <li>Guidebook for managers on breastfeeding, breastfeeding practices and different management models for breastfeeding in the workplace.</li> </ul>

ACAS, Advisory, Conciliation and Arbitration Service; CoC, Chain of Command; DOD, Department of Defense; HSE, Health and Safety Executive; MOD, Ministry of Defence; RAAF, Royal Australian Air Force; RAF, Royal Air Force; RTW, return to work; USAF, US Airforce.

long-term benefits for retention needed to be recognised.<sup>36 39 44</sup> Workplace attitudes can be addressed by changing organisational culture and adopting protective breastfeeding-specific policies.<sup>31 32 34 36–38 43</sup>

## **Facility provision**

Organisational policies for provision, <sup>30</sup> <sup>32</sup> <sup>34</sup> <sup>36</sup> <sup>38</sup> knowledge of and facility existence<sup>32</sup> <sup>36</sup> <sup>38</sup> are prerequisites for access. <sup>32</sup> <sup>34</sup> <sup>36</sup> However, without national legislation and sufficient research on RTW breastfeeding, policy development often stalls. <sup>31</sup> <sup>37</sup> All studies identified timely knowledge of workplace facility existence as vital in preventing mothers prematurely ceasing breastfeeding on false assumptions. This was most common in male-dominated organisations, <sup>30</sup> <sup>32</sup> <sup>35</sup> <sup>42</sup> <sup>43</sup> as was maternal concealment of breastfeeding, which prevented supervisors from facilitating appropriate access. <sup>22</sup> <sup>35</sup> <sup>38</sup> <sup>43</sup>

Restricted physical access, through limited opening times for mothers working non-standard hours,<sup>39 41 42</sup> or need of a key,<sup>34</sup> was a barrier. Advance warning of changes in working patterns, hours<sup>42</sup> or location<sup>37–39</sup> enabled arrangements for ongoing access to be made. Where employers offered only indirect breastfeeding facilities, mechanical,<sup>32 41</sup> psychological<sup>33 39 43 44</sup> and financial<sup>39</sup> difficulties transitioning from direct breastfeeding were frequently cited barriers. While an on-site nursery can facilitate access,<sup>32 34</sup> this is not always practical. Alternative off-site access<sup>32–34 37 40 42 43</sup> can be enabled by workplace accommodations, including shift pattern changes,<sup>33 42</sup> flexible<sup>32–34 41</sup> or parttime working,<sup>32–34 42</sup> or additional leave.<sup>33 40–43</sup>

## Use of facility

Facility suitability influenced maternal use. Synthesising these findings, we found that essential requirements were privacy and safety. <sup>30</sup> <sup>32</sup> <sup>34</sup> <sup>36</sup> <sup>39</sup> <sup>41-43</sup> Ideal facilities were designated and lockable, <sup>30</sup> <sup>32</sup> <sup>34</sup> <sup>36</sup> <sup>39</sup> <sup>41-43</sup> with basic amenities, including electricity and running water, <sup>32</sup> <sup>34</sup> <sup>36</sup> <sup>38</sup> <sup>41</sup> <sup>43</sup> even in austere military working environments. <sup>37</sup> <sup>38</sup> Optimal use of facilities<sup>34</sup> <sup>36-39</sup> <sup>43</sup> came with provision of refrigerated storage or equipment loans of hospital-grade pumps. <sup>34</sup> <sup>36-39</sup> <sup>43</sup> Unsuitable locations were toilets, public spaces and thoroughfares, <sup>32</sup> <sup>34-37</sup> <sup>39</sup> <sup>41</sup> <sup>43</sup> and those which exposed mothers to breastmilk transmissible hazards. <sup>22</sup> <sup>32</sup> <sup>38</sup> <sup>41</sup> Lack of advice or logistical support for transporting breastmilk during periods of separation resulted in breastfeeding cessation. <sup>38</sup>

Appropriate access was enabled through maternal antenatal<sup>35 36 44</sup> and universal workforce education,<sup>30-32 34-38 43</sup> which in three military studies was best received when delivered by HCPs.<sup>36 38 44</sup> Addressing maternal and supervisor expectations was equally important with use of facilities and time to breastfeed optimised through formally written and agreed breastfeeding plans.<sup>34 35 37-39 42 43</sup> Six studies<sup>32 36 37 39 41 43</sup> identified insufficient time as a barrier to accessing facilities. This was caused by occupational stress<sup>34 35 37 39 41 42</sup>; time-consuming, unpredictable workloads; inability to manage one's own schedule<sup>34 42</sup>; or limited flexible-working options.<sup>32-34 37 39 42</sup> Unplanned changes to workload,<sup>35 39</sup> working location<sup>39</sup> and patterns<sup>42</sup> all negatively impacted use of facilities.



Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram.<sup>27</sup> HIC, high-income country.

## DISCUSSION

Maternal motivation to breastfeed, which is predominantly driven by factors external to the workplace, was the most commonly identified factor enabling access of facilities.<sup>30 33–35 37–44</sup> The five key areas identified from policy analysis and the potential options for future policy development are explored and discussed based on the key barriers and enablers are identified. Recommendations focus on those factors which the British Army, as an employer, could influence.

## **Specific responsibilities**

Maternal duty to inform her employer<sup>18 19 23 47-59</sup> and medical officer<sup>47 48 52-54 59</sup> of breastfeeding status was the most commonly identified responsibility and is required in the UK.<sup>23</sup> However, numerous studies found actual or perceived lack of support,<sup>34 37 39 41 42</sup> pressure to stop or tolerance of hostile behaviours towards breastfeeding mothers<sup>30 41</sup> was a barrier to disclosure and accessing facilities. Non-disclosure could harm both maternal and infant health. The British Army has a duty

to protect the breastfeeding servicewoman and risk assess the potential impact of exposure to any of the known harmful occupational hazards identified in two observational studies and a comprehensive literature review.<sup>18 22</sup>

Disclosure, facility provision, access and use can be improved through breastfeeding-specific policy<sup>18</sup> <sup>19</sup> <sup>42</sup> <sup>51</sup> and universal workplace education.<sup>42</sup> <sup>54</sup> <sup>57</sup> Workplace breastfeeding policy is supported as good practice by Advisory, Conciliation and Arbitration Service<sup>19</sup> and recommended by the NHS,<sup>51</sup> HSE<sup>18</sup> and the US Army.<sup>42</sup> Following good practice from the USA, a multidisciplinary working group to evaluate policy implementation and breastfeeding outcome should be established.<sup>42</sup> However, policies must be appropriate<sup>38</sup> <sup>43</sup>; promoting extended maternity leave for breastfeeding fuels the misconception that it is unacceptable in the workplace.<sup>30</sup> <sup>40</sup> <sup>43</sup> To be effective, policies must empower the servicewomen, must be widely available<sup>30</sup> <sup>32</sup> <sup>34</sup> <sup>36</sup> and must be provided in a timely manner before RTW,<sup>32</sup> <sup>36</sup> <sup>38</sup> ideally in the antenatal period.<sup>32</sup> <sup>24</sup> <sup>36</sup> <sup>44</sup>

#### Table 5 Key areas in document

		Identifies specific maternal, employer, CoC and medical chain responsibilities	Identifies requirement for individualised breastfeeding plans or risk assessments	Details requirements and recommendations for breastfeeding facility access and provisions	Advises on provisions for postpartum physical fitness assessment	Signposts possible workplace accommodations for postpartum mothers
UK	HSE <sup>18</sup>	х	х	Х		
	ACAS <sup>19</sup>	х	х	Х		Х
	NHS <sup>51</sup>	Х	Х	Х		Х
HICs' Armed	MOD 59	Х	Х			Х
Forces	MOD <sup>43</sup>		Х	Х		
	RN <sup>60</sup>	Х			Х	
	RAF <sup>52</sup>	Х		Х	Х	
	RAF <sup>53</sup>	Х		Х		
	USAF <sup>54</sup>	Х	Х	Х		Х
	USAF 44				Х	Х
	DOD <sup>48</sup>	Х	Х	Х		Х
	DOD <sup>43</sup>	Х	Х	Х	Х	Х
	USCG <sup>55</sup>	Х	Х	Х		Х
	DOD 56	Х	Х	Х		Х
	DOD 57	х		Х		Х
	RAAF <sup>49</sup>	х	Х	Х	х	Х

ACAS, Advisory, Conciliation and Arbitration Service; CoC, Chain of Command; DOD, Department of Defense; HSE, Health and Safety Executive; MOD, Ministry of Defence; RAAF, Royal Australian Air Force; RAF, Royal Air Force; RN, Royal Navy; USAF, US Airforce; USCG, US Coast Guard.

Policy implementation is most successful when accompanied by universal workforce education, which tackles stigma and barriers and improves practical knowledge and awareness of the benefits of breastfeeding.<sup>30-32</sup> <sup>34-38</sup> <sup>43</sup> All US military studies which trialled educational programmes found military HCP delivery to have the greatest efficacy.<sup>36 38 44</sup> The US Army medical chain is now responsible for breastfeeding self-learning<sup>42</sup> and education delivery,<sup>42,54</sup> and any British Army programme would need to start with HCP education. HCP engagement was crucial in 60% of studies in advocating and supporting improved access and use of facilities.<sup>35–38 43 44</sup> While policy and education are commonly cited organisation responsibilities which demonstrate support and organisational acceptance, <sup>31 32 34 36-38 43</sup> alone they are insufficient; true change comes through creation of a supportive organisation culture where inappropriate behaviours are challenged.

#### Individualised plans and risk assessments

Three military studies<sup>32 36 38</sup> and most policies<sup>42 48 49 54-56</sup> identify line managers' and medical officers as respectively responsible for completing individualised risk assessments<sup>18 19 23 42 48 49 51 54-56</sup> and medical grading assessments,<sup>42 48 52 53 60</sup> ideally prior to RTW.<sup>19 23 42 48 55 56</sup> Line manager risk assessment should not, however, be replaced by medical grading.<sup>49 54 55</sup> In most cases, reasonable adjustments for breastfeeding should be an employer and not a medical responsibility; however, it may be appropriate to apply medical workplace restrictions due to individual postpartum functionality.<sup>22,59</sup> In line with HSE legislation,<sup>18</sup> triservice British Armed Forces policy does recommend individualised risk assessment.<sup>23,59</sup>

In addition to risk assessments, eight studies and six policies, including from the US Army and Royal Air Force, which provide the paperwork for completion, recommend individualised breastfeeding plans either concurrently or independently.<sup>35 37-39 43</sup> Well-considered and successfully implemented plans help manage expectations of the servicewomen, line managers and colleagues and counter misconceptions about workforce cohesion and operational capability.<sup>36 39</sup> Plans should be thorough, account for maternal breastfeeding practices and infant breastmilk requirements; notice for changes to work pattern or location<sup>37-39 42 47 48</sup>; and what facilities can be optimally provided in a range of settings, including the home base, courses, exercises and deployments.<sup>47 49</sup>

Although only identified by a few studies, mothers need time to psychologically prepare and practically plan adjustments to breastfeeding practices, storage and transportation. If cessation is required and timelines are insufficient, or facilities in a new location are inappropriate or inaccessible, medical

Table	6 Five key recommendations to enable access to breastfeeding on return to work in the British Army
1	Easily accessible breastfeeding-specific policy is required, optimised by online available guides.
2	A Women's Health Special Interest Group should ensure appropriate materials are available for <i>military HCP education</i> on breastfeeding in <i>Universal Workplace and Commanders Education</i> , and awareness is a responsibility of unit medical officers and personnel branches. Support and education for breastfeeding servicewomen particularly focusing on the challenges of military occupation and breastfeeding should be complemented by the development of a military-specific <i>servicewomen's breastfeeding network</i> .
3	Commanders and servicewomen must ensure <i>individual risk assessments and breastfeeding plans</i> are completed. Regular review as maternal and infant breastfeeding requirements change is also required. A breastfeeding model framework, as laid out by the RAAF, could be included in breastfeeding guides. This will manage maternal expectations and give line managers a framework within which to practically manage individual needs.
4	Breastfeeding policy must clarify minimum breastfeeding facility standards and those which should be aspired to. This should consider what is achievable in the home base, on courses, exercises and deployments. At an individual level, the breastfeeding plan should document expected provisions for each setting or workplace.
5	Employment policy should consider the option to extend exemptions from deployment, on an individual basis, to 24 months, allowing servicewomen to meet the WHO's 2-year breastfeeding target. Outcomes of ongoing postnatal research should be used to inform any recommendations on physical activity or fitness testing.

HCP, healthcare professional; RAAF, Royal Australian Air Force.

complications such as engorgement, pain or mastitis<sup>50</sup> can occur and a period of absence and treatment may be required. Compromise and circumstantially appropriate plans should be agreed, balancing the needs of the service; the servicewoman's career; and her practical, psychological and medical breastfeeding needs.<sup>34 35 37-39 42 43</sup>

#### **Breastfeeding facilities**

For mothers to access facilities, they must exist.<sup>33</sup> <sup>34</sup> <sup>36</sup> <sup>43</sup> Most policies and studies<sup>30</sup> <sup>32</sup> <sup>34</sup> <sup>36</sup> <sup>39</sup> <sup>41-43</sup> identify privacy and safety as the minimum standard, even for exercises and deployments.<sup>47</sup> This, along with space to rest, is legally required in UK workplaces.<sup>18</sup> Toilets, public thoroughfares and locations with hazardous exposures were universally recognised as unsuitable.<sup>18</sup> <sup>19</sup> <sup>22</sup> <sup>23</sup> <sup>34</sup> <sup>37</sup> <sup>39</sup> <sup>41</sup> <sup>42</sup> <sup>47-49</sup> <sup>51-57</sup> US and Australian policies specify minimum requirements for exercises and deployments,<sup>47</sup> and recommend access to running water and electricity.<sup>47-49</sup> <sup>54-58</sup> Water is required for hygiene reasons and to prevent infection.<sup>7 9</sup> <sup>16</sup> <sup>17</sup> <sup>20</sup> <sup>23</sup> Electricity is a necessity for electric pumps, but manual pumps and hand expression could be considered; it is also ideal for refrigerated storage; however, cold storage boxes can be temporarily used.

Refrigerated storage was recommended in the UK,<sup>19 51</sup> Australia<sup>49</sup> and the USA for shifts over 12 hours.<sup>55</sup> Although UK workplace legislation is lacking,<sup>15</sup> Ministry of Defence (MOD) policy does make provisions for privacy and breastmilk storage,<sup>23 53 59</sup> and this is identified as an enabler. In other countries, cold-storage provision is a maternal responsibility, although support with arranging cold chains for transportation from deployments or austere environments is an enabler.<sup>19 47 48 54</sup> Provision of pumping equipment was consistently a maternal responsibility; however, access to hospital-grade loan pumps was viewed as beneficial by servicewomen and employers.<sup>47 48 5</sup> Firm-based locations<sup>23</sup> should provide electricity and refrigerated storage. To enable appropriate planning.<sup>37-39</sup> locations and circumstances where these cannot be provided, alongside the essentials of privacy, safety and running water, should be made clear.

Sufficient time to access facilities should be provided, <sup>19</sup> <sup>47</sup> <sup>49</sup> <sup>51</sup> <sup>54</sup>-<sup>60</sup> determined by maternal needs, <sup>49</sup> <sup>51</sup> <sup>56</sup> individual breastfeeding practices <sup>49</sup> or medical recommendations. <sup>56</sup> Facility provision is a futile gesture if mothers cannot access them. For mothers unable to manage their own schedule, <sup>34</sup> <sup>37</sup> <sup>39</sup>-<sup>43</sup> formally agreed breastfeeding plans are the most commonly identified enabler. <sup>33</sup> <sup>36</sup> <sup>41</sup> <sup>43</sup> RAAF breastfeeding models <sup>49</sup> provide guidance and frameworks to manage expectations of break duration and frequency.

#### **Physical fitness provisions**

Two studies identify fear of career ramifications, secondary to failure to meet physical fitness requirements or to attend career courses, as a barrier to accessing workplace facilities.<sup>39 41</sup> However, challenges of physical fitness are not just restricted to breastfeeding mothers and are considered in various nations' policies on military RTW physical fitness provisions for all postpartum mothers.<sup>48 49 52 58 60</sup> These include providing specialist postnatal Physical Training (PT),<sup>49 52 60</sup> deconflicting breastfeeding breaks with scheduled PT,<sup>48</sup> and exempting physical testing for 90 days and 6<sup>58</sup> or 12 months.<sup>52</sup> British Army policy makes provisions for this; however, future recommendations will be reviewed, in line with results from ongoing postpartum musculoskeletal studies.

## Systematic review

## Workplace accommodations

Fear or actual infant separation can cause practical storage and transportation and psychological difficulties when transitioning from direct breastfeeding to expressing.<sup>33 39 43 44</sup> Longer and uncertain duration often results in breastfeeding cessation. Separation from a formed breastfeeding support network is also a barrier to continuation of breastfeeding, although many military studies found a virtual military-specific peer network to be enabling.<sup>35 39 41 43 44</sup> To protect from early separation, all HICs' Armed Forces recommend exemption from deployment,<sup>47 49 54–59</sup> but duration varies from 6 to 24 months postpartum. 47 49 54-59 The British Army offer 6 months, with provision to extend to 12 months at the commanders discretion.<sup>59</sup> Numerous studies and policies<sup>47 49 51</sup> recognise the beneficial impact flexible-working and part-time working policies can have on maternal access of appropriate facilities.<sup>32 33 37 40 43</sup> Such workplace accommodation policies are now offered subject to the needs of the service, but should be signposted in breastfeeding policy.

## **Recommendations**

Five key recommendations were drawn from this work (Table 6), and these have been presented to the Director of Personnel department, Women's Health Advisory Group and Primary Care Women's Health Special Interest Group.

## CONCLUSIONS

RTW breastfeeding benefits the infant, mother and employer. However, numerous factors can either enable or present a barrier to breastfeeding. Thematic analysis identified three key areas impacting access: attitudes towards breastfeeding, issues relating to facility provision and those related to use. Dominant workplace themes influencing access included breastfeeding policy protecting mothers and facilities, workplace support to access facilities, universal workforce education, provision of suitable breastfeeding facilities, individualised risk assessments and breastfeeding plans, and planning for infant and breastfeeding support network separation. Five key recommendations were made when these themes were considered in relation to British Army employment. There remains a need for further research and appropriate monitoring, evaluation and oversight of British Army breastfeeding implementation to inform subsequent policy amendments.

Contributors This research and article was undertaken by HT as sole author.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

**Data availability statement** Data sharing is not applicable as no datasets were generated and/or analysed for this study. There are no datasets used or produced in this study.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

#### ORCID iD

Hannah Taylor http://orcid.org/0000-0002-9781-2085

## Systematic review

#### REFERENCES

- 1 Kramer MS, Kakuma R. *The optimal duration of exclusive breastfeeding: a systematic review*. Geneva: WHO, 2001.
- 2 NHS. Benefits of breastfeeding, 2017. Available: http://www.nhs.uk/Conditions/ pregnancy-and-baby/Pages/benefits-breastfeeding.aspx [Accessed 17 Jul 2017].
- 3 Horta BL, Loret de Mola C, Victora CG. Breastfeeding and intelligence: a systematic review and meta-analysis. Acta Paediatr 2015;104:14–19.
- 4 Dieterich CM, Felice JP, O'Sullivan E, et al. Breastfeeding and health outcomes for the mother-infant dyad. *Pediatr Clin North Am* 2013;60:31–48.
- 5 Chowdhury R, Sinha B, Sankar MJ, et al. Breastfeeding and maternal health outcomes: a systematic review and meta-analysis. Acta Paediatr 2015;104:96–113.
- 6 Hahn-Holbrook J, Schetter CD, Haselton M. Breastfeeding and maternal mental and physical health. In: Spiers MV, Geller PA, Kloss JD, eds. Women's health psychology. Wiley: Hoboken NJ, 2013.
- 7 Cohen R, Mrtek MB, Mrtek RG. Comparison of maternal absenteeism and infant illness rates among breast-feeding and formula-feeding women in two corporations. *Am J Health Promot* 1995;10:148–53.
- 8 8Del Bono E, Pronzato CD. Does breastfeeding support at work help mothers and employers at the same time? Bonn: Institute for the Study of Labor (IZA), 2012.
- 9 Ortiz J, McGilligan K, Kelly P. Duration of breast milk expression among working mothers enrolled in an employer-sponsored lactation program. *Pediatr Nurs* 2004;30:111–9.
- 10 Oakley LL, Renfrew MJ, Kurinczuk JJ, et al. Factors associated with breastfeeding in England: an analysis by primary care trust. *BMJ Open* 2013;3. doi:10.1136/ bmjopen-2013-002765. [Epub ahead of print: 20 Jun 2013].
- 11 Spencer RL, Greatrex-White S, Fraser DM. 'I thought it would keep them all quiet'. Women's experiences of breastfeeding as illusions of compliance: an interpretive phenomenological study. J Adv Nurs 2015;71:1076–86.
- 12 NHS. Infant feeding survey 2010. The NHS Information Centre: New York, 2012.
- 13 NICE. Promotion of breastfeeding initiation and duration. London: NHS, 2006.
- 14 Victora CG, Bahl R, Barros AJD, et al. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. Lancet 2016;387:475–90.
- 15 Gray Het al. World breastfeeding trends initiative: UK report 2016. London: The World Breastfeeding Trends Initiative, 2016.
- 16 World Policy Centre. Are mothers of infants guaranteed breastfeeding breaks at work? 2017. Available: https://www.worldpolicycenter.org/policies/are-mothers-of-infantsguaranteed-breastfeeding-breaks-at-work [Accessed 2 Jul 2017].
- 17 Health and Safety Commission. The workplace (health, safety and welfare) regulations 1992. provision 25 (4). Available: http://www.legislation.gov.uk/uksi/1992/3004/ regulation/25/made [Accessed 23 Jun 2017].
- 18 HSE. New and expectant mothers who work: a brief guide to your health and safety, 2013. Available: http://www.hse.gov.uk/pubns/indg373.pdf [Accessed 7 Aug 2017].
- 19 ACAS. Accommodating breastfeeding employees in the workplace, 2013. Available: http://www.acas.org.uk/media/pdf/2/i/Acas-guide-on-accommodating-breastfeedingin-the-workplace.pdf [Accessed 12 Jul 2017].
- 20 Baker M, Milligan K. Maternal employment, breastfeeding, and health: evidence from maternity leave mandates. J Health Econ 2008;27:871–87.
- 21 MOD. Adaptability and partnership: issues for the strategic defence review. London: The Stationery Office, 2010.
- 22 Croft AM. The employability of pregnant and breastfeeding servicewomen. J R Army Med Corps 1995;141:134–41.
- 23 MOD. Joint service publication 375: management of health and safety in defence. London: MOD, 2017.
- 24 Li R, Fein SB, Chen J, et al. Why Mothers Stop Breastfeeding: Mothers' Self-reported Reasons for Stopping During the First Year. *Pediatrics* 2008;122:S69–76.
- 25 Defence Committee. SDSR 2015 and the army. 2017. London: House of Commons, 2015.
- 26 MOD. UK armed forces biannual diversity statistics. London: Office National Statistics, 2016.
- 27 27Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009;6:e1000097.
- 28 Critical Appraisal Skills Programme. Casp Checklisst, 2017. Available: https://casp-uk. net/casp-tools-checklists/ [Accessed 6 Aug 2017].
- 29 Strengthening the reporting of observational studies in epidemiology. STROBE crosssectional studies checklist, 2017. Available: https://strobe-statement.org/fileadmin/ Strobe/uploads/checklists/STROBE\_checklist\_v4\_cross-sectional.pdf [Accessed 6 Aug 2017].
- 30 Gatrell CJ. Secrets and lies: breastfeeding and professional paid work. Soc Sci Med 2007;65:393–404.

- 31 Hawkins SS, Griffiths LJ, Dezateux C, et al. The impact of maternal employment on breast-feeding duration in the UK millennium cohort study. *Public Health Nutr* 2007;10:891–6.
- 32 32Kosmala-Anderson J, Wallace LM. Breastfeeding works: the role of employers in supporting women who wish to breastfeed and work in four organizations in England. *Public Health Nutrition* 2007;10:891–6.
- 33 Skafida V. Juggling work and motherhood: the impact of employment and maternity leave on breastfeeding duration: a survival analysis on growing up in Scotland data. *Matern Child Health J* 2012;16:519–27.
- 34 Wallace LM, Kosmala-Anderson J, Mills S, et al. Mutually exclusive? A United Kingdom survey of women's experiences of breastfeeding and working. *Midwifery Digest* 2008;18:99–103.
- 35 Bales K, Washburn J, Bales J. Breastfeeding rates and factors related to cessation in a military population. *Breastfeed Med* 2012;7:436–41.
- 36 Bell NR, Ritchie EC. Breastfeeding in the military: Part I. information and resources provided to service women. *Mil Med* 2003;168:807–12.
- 37 Martin SE, Drake E, Yoder L, et al. Active duty women's perceptions of breast-feeding support in the military setting. *Mil Med* 2015;180:1154–60.
- 38 Sleutel MR. Breastfeeding during military deployment: a soldier's story. *Nursing for Women's Health* 2012;16:20–5.
- 39 Stevens KV, Janke J. Breastfeeding experiences of active duty military women. *Mil Med* 2003;168:380–4.
- 40 Stewart K. Military milk: breastfeeding rates among Australian defence force women who return to military service following maternity leave. J Hum Lact 2015;31:138–44.
- 41 Uriell Z, Perry A, Kee A, et al. Breastfeeding in the navy: estimates of rate, duration, and perceived support. *Mil Med* 2009;174:290–6.
- 42 Zilanawala A. Maternal nonstandard work schedules and breastfeeding behaviors. Matern Child Health J 2017;21:1308–17.
- 43 Bristow KM. Barriers and facilitators of breastfeeding for primiparous active duty military mothers: a qualitative study. Ipswich, MA: Uniformed Services University of the Health Sciences, 1999.
- 44 Harlow BA. Breastfeeding counselling, barriers and facilitators of lactation in the military community. Ipswich, MA: Uniformed Services University of the Health Sciences, 1998.
- 45 Jacobs B, Ir P, Bigdeli M, et al. Addressing access barriers to health services: an analytical framework for selecting appropriate interventions in low-income Asian countries. *Health Policy Plan* 2012;27:288–300.
- 46 Kelle U. Combining qualitative and quantitative methods in research practice: purposes and advantages. *Qualitative Research in Psychology* 2006;3:293–311.
- 47 Department of the Army. Army directive 2015-43: memorandum for the revised breastfeeding and lactation support policy. Washington: Department of Defense, 2016.
- 48 Department of the Army. *Memorandum for breastfeeding support plan on return to work*. Washington: Department of Defense, 2016.
- 49 The Airforce.. *Air force diversity Handbook: guidance for Commanders and managers of breastfeeding mothers*. Canberra: The Royal Australian Air Force, 2014.
- 50 La Leche League GB. When breastfeeding ends suddenly, 2017. Available: https:// www.laleche.org.uk/breastfeeding-ends-suddenly/ [Accessed 3 Sep 2017].
- 51 NHS. Breastfeeding and work: information for employees and employers. London: Department of Health, 2008.
- 52 The Royal Airforce.. Pregnancy in the RAF: a guide for servicewomen. London: The Royal Airforce, 2014.
- 53 The Royal Airforce.. Pregnancy in the RAF: a guide for line managers. London: The Royal Airforce, 2017.
- 54 Department of the Airforce. Air force instruction 44-102: medical care management. Washington: Department of Defense, 2017.
- 55 United States Coast Guard. *Coast guard COMDINST M1000.9: pregnancy policy and guidelines*. Washington: US Department of Homeland Security, 2016.
- 56 Department of the Navy. Marine Corps order 5000.12E: marine Corps policy concerning pregnancy and parenthood. Washington: Department of Defense, 2004.
- 57 Department of the Navy. OPNAV instruction 6000: 1C: US navy policy concerning pregnancy and parenthood. Washington: Department of Defence, 2007.
- 58 Department of the Airforce. Us air force memorandum on post-pregnancy deferment and exemption periods. Washington: Department of Defense, 2015.
- 59 MOD. Joint service publication 760: tri-service regulations for leave and other types of absences. London: MOD, 2016.
- 60 The Royal Navy. BR3(1): naval personnel management. Portsmouth: The Royal Navy, 2016.