



OPEN ACCESS

Degenerative cervical spine changes among early career fighter pilots: a 5-year follow-up

Tuomas Keskimölo,¹ J Pernu,¹ J Karppinen,^{2,3} J Niinimäki,^{1,2} P Oura,² T Leino,^{4,5} T Honkanen⁶

¹Faculty of Medicine, University of Oulu, Oulu, Finland

²Medical Research Center Oulu, Oulu University Hospital, Oulu, Finland

³Finnish Institute of Occupational Health Oulu Regional Office, Oulu, Finland

⁴Air Force Command Finland, Tikkakoski, Finland

⁵Department of Leadership and Military Pedagogy, National Defence University, Helsinki, Finland

⁶Aeromedical Centre, Centre for Military Medicine, Finnish Defence Forces, Helsinki, Uusimaa, Finland

Correspondence to

Tuomas Keskimölo, Faculty of Medicine, University of Oulu, Oulu, Finland; tuomas.keskimolo@student.oulu.fi

Received 24 March 2021

Accepted 29 May 2021

ABSTRACT

Introduction Degenerative changes of the cervical spine often cause disability and flight duty limitations among Finnish Air Force (FINAF) fighter pilots. We aimed to study the effect of +Gz exposure on degenerative changes in the cervical spine by comparing cervical MRIs of FINAF fighter pilots and controls.

Methods At baseline, the volunteer study population consisted of 56 20-year-old FINAF male fighter pilots (exposure group) and 56 21-year-old Army and Navy cadets (control group). Both groups underwent MRI of the cervical spine at the baseline and after 5 years. Degenerative changes evaluated using MRI included intervertebral disc (IVD) degeneration (Pfirrmann classification), disc herniations, uncovertebral arthrosis, Schmorl's nodes, Modic changes, spinal canal stenosis, kyphosis and scoliosis.

Results The degree of IVD degeneration in the whole cervical spine increased significantly in both populations with no between-group differences. The prevalence of disc herniations also tended to increase in both populations with no difference in the incidence over the follow-up. However, pilots proved to have more disc herniations at the baseline and at the follow-up. There were virtually no between-group differences in other assessed degenerative changes.

Discussion We found that IVD degeneration and the prevalence of disc herniations increased at a similar rate for fighter pilots and non-flying military students when all cervical levels were summed up. The lack of difference may be explained by the relatively low cumulative +Gz exposure during the first 5 years of a pilots' career.

INTRODUCTION

Fighter pilots need to move their heads with full range of motion in three directions to observe the airspace around them with simultaneous exposure to high +Gz forces.¹ The combination of +Gz forces and awkward head positions is potentially harmful for the structures of the cervical spine.² Neck pain is reported to be a common health problem among military pilots.^{3,4} A meta-analysis, consisting of 18 studies and more than 8000 pilots, revealed that 51% of fighter pilots suffered from neck disorders, and one-third of them had sought medical care for neck pain.⁴

Degenerative changes of the cervical spine in MRI include decreased signal intensity and height of intervertebral disc (IVD), annular fissures, protrusions/herniations and bone marrow lesions.⁵ In a meta-analysis by Shiri *et al*,⁶ neck pain was more common among pilots exposed to high

Key messages

- ▶ Fighter pilots are subject to high +Gz forces and awkward neck positions while flying.
- ▶ It has been proven that the load-bearing capacity of the neck muscles can be exceeded during combat manoeuvring.
- ▶ This may be harmful for the structures of the cervical spine, potentially leading to premature degeneration.
- ▶ The present study suggests that cervical spine degeneration is present in both flying and non-flying young healthy military male personnel.
- ▶ The accessory effect of G-force stress in premature cervical spine degeneration is yet to be reliably discovered.

G-forces compared with pilots flying less G-capable aircraft, but the authors found no difference between the groups in terms of cervical IVD degeneration. However, the results of individual studies on cervical IVD degeneration are conflicting. For example, Hämäläinen and Petren-Mallmin and Linder found that there was an increased prevalence of degenerative changes in MRI among fighter pilots when compared with age-matched controls.^{7,8} Petren-Mallmin and Linder⁸ found that experienced military pilots had significantly more osteophytes, protrusions/herniations and both spinal and foraminal stenosis. Furthermore, Hendriksen and Holewijn⁹ compared F-16 pilots to pilots flying less G-capable aircraft and found that frequent exposure to high +Gz forces may cause premature degenerative changes. However, several other studies¹⁰⁻¹² did not observe a significant difference in cervical IVD degeneration between the pilots and controls.

The contributing factors to fighter pilots' IVD degeneration are suggested to be the head movements during the Gz exposure and cumulative trauma thereof. A recent study investigated in-flight neck muscle activity during combat manoeuvring. The results implicated that high +Gz combined with awkward head postures (eg, rotations in combination with extensions) causes a force exceeding the load-bearing capacity of the cervical muscles.¹ This possibly results in more loading to the bony structures, IVDs and ligaments of the spine and may be a risk factor for early IVD degeneration. Furthermore, it has been suggested that the cumulative soft tissue (muscle or ligament) injuries may lead to



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Keskimölo T, Pernu J, Karppinen J, *et al*. *BMJ Mil Health* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bmjilitary-2021-001848

unstable spinal structures that lead to abnormal IVD loads and later to degeneration.

Neck disorders may lead to temporary or permanent flight disqualifications or limitations.^{13 14} In the Finnish Air Forces (FINAF), disability accompanied by spinal IVD degeneration is reported as the most common reason for aeromedical limitations.³ These flight duty limitations do not only cause individual harm but are economically and operationally significant. Fighter pilot training is highly expensive, which limits the number of pilots selected for fighter training. In order to preserve a sufficient number of fighter pilots, the pilots must maintain their working capability both physically and mentally. Therefore, it is important to take care of and monitor the health of fighter pilots already in the training phases.

The primary aim of the present study was to investigate the effect of +Gz exposure on degenerative changes of the cervical spine by comparing FINAF fighter pilots and controls. The degenerative changes of the cervical spine (from C2 to Th1) were assessed by MRI.

METHODS

Subjects

At baseline, the volunteer study population consisted of 56 FINAF male fighter pilot cadets and 56 sex-matched Finnish Defense Force Army and Navy cadets from the National Defense University as controls. FINAF cadets were flying BAe Hawk jet trainer (up to +8 Gz) during the follow-up. Whereas the control group's occupations during follow-up varied mainly from office work to conscript trainers that do not expose to accessory neck strain or G force stress. All pilots and controls completed follow-up. A questionnaire considering health and physical condition was filled by both groups. The questionnaire included questions on age, weight, height and physical performance. This structured questionnaire has been used previously.¹⁵ We used the STROBE cohort reporting guidelines for this study (online supplemental file 1).

The results of the first Cooper's test (12 min maximal distance running test) as a conscript were used to determine the physical condition of the subjects. Before starting the research, the procedures were explained to all the participants, who gave their voluntary written informed consent under the Declaration of Helsinki.

MRI examination

The MRI examinations were performed at the baseline and after a 5-year follow-up for both populations, between 2008 and 2020. MRI images were acquired using clinical 1.5 T MRI scanners. The imaging protocols included conventional T1-weighted and T2-weighted fast spin echo sequences: for example, 11 ms TE (Echo Time) and 440 ms TR (Repetition Time) for T1-weighted and 113 ms TE and 3300 ms TR for T2-weighted sagittal images and 100 ms TE and 3700 ms TR for T2-weighted axial images. The slice thickness was 3 mm, and the in-plane resolution was 0.47×0.47 mm for the sagittal plane and 0.35×0.35 mm for the axial plane.

The degree of IVD degeneration was assessed using the Pfirrmann classification with five grades consisting of evaluations of the disc space height, disc homogeneity, nucleus intensity and distinction between nucleus and annulus.¹⁶ An IVD degeneration sum score was composed by adding the scores of degenerated discs (Pfirrmann grades 1 and 2=0p, grade 3 (mild degeneration)=1p, grade 4 (moderate degeneration)=2p, grade 5 (severe degeneration)=3p) together and divided by the number

of subjects.¹⁷ IVD herniations were classified as protrusions, extrusions and sequesters but were later combined due to the low number of extrusions and sequesters.¹⁸ Schmorl's nodes were evaluated from each vertebral body throughout the cervical spine.¹⁹ Osteophytes were assessed from both sides of vertebral bodies and classified as uncovertebral arthrosis whether the osteophytes were unilateral or bilateral.²⁰ Scoliosis was noted if a scoliotic angle of at least 10° was measured.²¹ The spinal rotation was not measured as we did not address a medical diagnosis of scoliosis. The kyphotic angle was measured using the Harrison posterior tangent method and recorded if the kyphotic angle in the cervical spine was at least 1°.²² Spinal canal stenosis was first recorded at each cervical level individually but was later simplified to comprise the whole cervical spine.²³ Modic changes (types 1, 2 and 3) were evaluated using both T1-weighted and T2-weighted images as described.²⁴

The MRI scans were read by two persons including a musculoskeletal radiologist. The scans were not assessed blinded due to data security practices concerning pilot information. Both examiners were blinded to each other's findings. The intrarater and inter-rater reliabilities were assessed using kappa and weighted kappa coefficients.

Statistical analysis

Data from the MRI examinations and questionnaire were analysed using SPSS Statistics V.25 software for Windows. Descriptive statistics were presented as means and SDs for continuous variables with normal distributions, medians with IQRs for continuous variables with skewed distributions or percentages and frequencies (n) for categorical variables (exceptions are mentioned in table descriptions). Normality was evaluated by visual inspection. The tests used to determine the association between groups, symptoms and MRI findings included independent sample t-test for continuous variables with normal distributions, a Mann-Whitney test for continuous variables with skewed distributions, a χ^2 test for categorical variables, a Wilcoxon test for repeated measurements for continuous variables with skewed distributions and a McNemar test for repeated measurements for class variables. The threshold for statistical significance was set at $p=0.05$.

RESULTS

Subjects

There was no significant difference at the baseline in height, weight, BMI or last school grade of physical education between the groups. However, the pilots were younger and had performed better in their first Coopers test as a conscript (Table 1).

Table 1 Main characteristics of the samples at the baseline

Characteristic	Pilots (n=56)	Controls (n=56)	P for difference
Age (years)	20 (±0.7)	21 (±1.2)	<0.001
Height (cm)	179.9 (±4.7)	180.4 (±5.6)	0.650
Weight (kg)	75.7 (±6.3)	77.0 (±7.7)	0.346
BMI (kg/m)	23.4 (±1.6)	23.6 (±1.9)	0.467
Cooper's test* (m)	3040 (±190)	2780 (±420)	<0.001
Last physical education grade	9.29 (±0.62)	9.13 (±0.85)	0.258

The values are means with SD.
BMI, body mass index.

IVD degeneration

During the follow-up, a statistically significant progression in IVD degeneration was found in both groups. Progression was noted for both pilots and controls in IVD degeneration at all levels from C2/C3 to C6/C7 (Table 2) and in the IVD degeneration sum score (Table 3). At C7/Th1, the change was not significant in either group. When comparing the groups, the change in Pfirrmann score among the fighter pilots was of greater magnitude at C6/C7 ($p=0.007$). There was no significant difference in the change of the IVD degeneration sum score between the groups. There was no between-group difference at the baseline or follow-up (Table 3, Figure 1).

IVD herniations

The total number of IVD herniations increased significantly among controls ($p=0.004$) and showed an almost increasing trend among pilots ($p=0.061$). Both at baseline and follow-up, the fighter pilots had a higher prevalence of IVD herniations, but there was no difference in the incidence of IVD herniations between the groups (Table 3).

Other cervical degenerative changes

The prevalence of uncovertebral arthrosis (number of arthritic vertebral levels per individual) did not differ at baseline, but at follow-up, it was higher among the pilots. There was no difference in the incidence of uncovertebral arthrosis between the groups. No between-group differences were observed in Schmorl's nodes, Modic changes, kyphosis, scoliosis or stenosis at the baseline or follow-up. Two pilots developed type 1 Modic change during the follow-up, whereas one control had a type 2 Modic change at the baseline and developed one more type 2 Modic change during the follow-up (Table 3).

Rater reliability

All intrarater reliabilities were substantial or perfect (0.6–1.0). Inter-rater reliabilities were between 0.6 and 1.0 for disc herniations, Schmorl's nodes, Modic changes, kyphosis and scoliosis. There was more disagreement in uncovertebral arthrosis (0.3) and IVD degeneration (0.2) in which the disagreement was linear in nature.

DISCUSSION

According to the present findings, there was a progression of IVD degeneration in both groups. Additionally, the prevalence of disc herniations increased significantly among the controls and had an increasing trend of borderline significance among pilots. However, a more prominent progression in the fighter pilots with cumulative +Gz exposure was observed only in IVD degeneration at C6/C7. We did not find any progression of uncovertebral arthrosis, Schmorl's nodes, Modic changes, kyphosis, stenosis or scoliosis during the follow-up. We found higher prevalence of herniations in pilots at baseline and at follow-up. Because either of the populations had been exposed to G force stress at baseline and the other underlying factors are unknown, our primary aim was to assess the incidences. Also, the controls seem to trend worse in disc degeneration sum score medians at baseline (Table 3). However, the difference was not significant either in disc degeneration sum score medians or in average Pfirrmann scores (pilots 2.22, controls 2.35).

Control population selection

The control population proved to be physically very similar to the study population. There was no statistically significant

difference in height, weight, BMI or physical education grade between the groups. The difference in age proved to be statistically significant, but the difference was only 0.7 years. The notable difference in the average result of the Cooper's test may be partly explained by more demanding physical performance requirements during the application phase for the fighter pilot training.

Comparison to previous literature

The present findings of IVD degeneration were similar to the meta-analysis by Shiri *et al.*⁶ However, when comparing our results with individual studies, there were some conflicting results,^{7,8} whereas others were well in line with our results.^{10,11} When comparing the present results to previous studies, it has to be kept in mind that the previous investigations have had limitations such as a small number of subjects. Both Hämäläinen and Sovelius used only 12 fighter pilots with 12 age-matched controls.^{7,11} In Sovelius's study, degenerative changes in the cervical spine of fighter pilots were more common in the lower part of the cervical spine, especially at the C5/C6 and C6/C7 levels.¹¹ Instead, the degenerative changes in the control group were more scattered in both studies.^{7,11}

The fighter pilots had a total experience of 5 years of military flying and only 3 years of flying with high performance aircraft (up to +8Gz). A comparison between our study and other studies must be made with caution, since the Hawk flight hours were below 200.

Magnetic resonance imaging

MRI is widely used in this research field. The Harrison posterior tangent method is an accurate method for estimating the curvature of the cervical spine, and it has lower SEs of measurement than the four-line or two-line Cobb methods.²² However, our intention was not to diagnose kyphosis but only to assess kyphotic angles in the images. MRI also shows the shape of IVDs and vertebrae clearly and is commonly used for recording Schmorl's nodes.¹⁹

The prevalence of some MRI findings (eg, Modic changes and spinal canal stenosis) was low, excluding any prominent differences between the groups. The inter-rater disagreement in IVD degeneration is endorsed by the fact that the Pfirrmann classification is primarily structured for and used in the assessment of lumbar IVDs, but it has been adapted also in assessment of cervical spine.^{16,25} Our aim was to assess the incidences between groups and not primarily the prevalences.

When comparing results with previous studies, it must be taken into account that they have used rather low-quality MRI devices. Hämäläinen used a 0.1 T MRI device, and Petren-Mallmin and Linder used a 0.5 T device throughout the study.^{7,8} Sovelius used a 0.1 T MRI device in the baseline phase, which was replaced by a 1.0 T device in follow-up phase.¹¹ In the Hendriksen and Holeyijn⁹ study, the mean age of the F-16 group was higher than the control group. After the groups were adjusted for age, the difference between the groups disappeared.

When comparing IVD degeneration of our population to civilian 21–30-year-old male population, the results were similar with Pfirrmann grades 2 and 3 being clearly the most common.²⁵ Our finding of the highest prevalence of disc herniations and IVD degeneration in the lower cervical spine (ie, C5/C6) is also similar to other studies. The lower cervical spine was most affected in non-flying 12–95-year-old population and age-matched FINAF fighter pilots.^{11,25}

Table 2 Prevalence of MRI findings among pilots and controls

Finding	Baseline							Follow-up						
	C2/3	C3/4	C4/5	C5/6	C6/7	C7/Th1	C2/3	C3/4	C4/5	C5/6	C6/7	C7/Th1		
Pilots														
Pfirrmann grade														
≤2	69.6 (39)	48.2 (27)	66.1 (37)	60.7 (34)	89.3 (50)	92.9 (52)	48.2 (27)	32.1 (18)	35.7 (20)	28.6 (16)	53.6 (30)	87.5 (49)		
3	30.4 (17)	51.8 (29)	33.9 (19)	37.5 (21)	10.7 (6)	5.4 (3)	51.8 (29)	66.1 (37)	62.5 (35)	55.4 (31)	39.3 (22)	12.5 (7)		
≥4	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	0.0 (0)	1.8 (1)	0.0 (0)	1.8 (1)	1.8 (1)	16.1 (9)	7.1 (4)	0.0 (0)		
Disc herniation	1.8 (1)	3.6 (2)	3.6 (2)	5.4 (3)	8.9 (5)	0.0 (0)	0.0 (0)	7.1 (4)	3.6 (2)	26.8 (15)	7.1 (4)	0.0 (0)		
Uncovertebral arthrosis	1.8 (1)	8.9 (5)	1.8 (1)	7.1 (4)	5.4 (3)	1.8 (1)	1.8 (1)	14.3 (8)	3.6 (2)	8.9 (5)	5.4 (3)	0.0 (0)		
Schmorl's node	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	0.0 (0)	3.6 (2)	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	0.0 (0)	1.8 (1)		
Modic change	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	1.8 (1)	0.0 (0)		
Spinal canal stenosis	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	7.1 (4)	1.8 (1)	0.0 (0)		
Kyphosis	30.4 (17)						21.4 (12)							
Scoliosis	3.6 (2)						3.6 (2)							
Controls														
Pfirrmann grade														
≤2	66.1 (37)	58.9 (33)	41.1 (23)	44.6 (25)	64.3 (36)	83.9 (47)	45.5 (25)	34.5 (19)	38.2 (21)	27.3 (15)	50.9 (28)	76.4 (42)		
3	33.9 (19)	33.9 (19)	41.1 (23)	46.4 (26)	32.1 (18)	16.1 (9)	50.9 (28)	54.5 (30)	54.5 (30)	56.4 (31)	45.5 (25)	23.6 (13)		
≥4	0.0 (0)	7.1 (4)	0.0 (0)	8.9 (5)	3.6 (2)	0.0 (0)	3.6 (2)	10.9 (6)	7.3 (4)	16.4 (9)	3.6 (2)	0.0 (0)		
Disc herniation	0.0 (0)	0.0 (0)	1.8 (1)	3.6 (2)	0.0 (0)	0.0 (0)	1.8 (1)	0.0 (0)	3.6 (2)	12.5 (7)	7.1 (4)	0.0 (0)		
Uncovertebral arthrosis	0.0 (0)	3.6 (2)	3.6 (2)	8.9 (5)	7.1 (4)	0.0 (0)	0.0 (0)	1.8 (1)	3.6 (2)	5.5 (3)	7.3 (4)	0.0 (0)		
Schmorl's node	1.8 (1)	1.8 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	5.4 (3)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)		
Modic change	0.0 (0)	0.0 (0)	1.8 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	1.8 (1)	0.0 (0)	0.0 (0)		
Spinal canal stenosis	0.0 (0)	0.0 (0)	0.0 (0)	3.6 (2)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	1.8 (1)	3.6 (2)	0.0 (0)	0.0 (0)		
Kyphosis	30.4 (17)						23.2 (13)							
Scoliosis	3.6 (2)						1.8 (1)							

The values are percentages (number of findings per individual) with frequencies (number of findings at each cervical level).

Table 3 Difference between pilots and controls in MRI findings

Finding	Pilots			Controls			Between-group differences		
	Baseline	Follow-up	P for change within group	Baseline	Follow-up	P for change within group	P for difference in baseline value between groups	P for difference in follow-up value between groups	P for difference in changes between groups
Disc degeneration sum score; median (IQR)	1.5 (1–3)	3 (2–5)	<0.001	3.0 (1–3.75)	4 (2–5)	<0.001	0.090	0.525	0.225
Disc herniation	23 (13)	45 (25)	0.061	5 (3)	25 (14)	0.004	0.015	0.020	0.689
Uncovertebral arthrosis	26.8 (15)	33.9 (19)	0.454	23.2 (13)	17.6 (10)	0.625	0.440	0.024	0.182
Schmorl's node	5.4 (3)	3.6 (2)	>0.999	3.6 (2)	7.1 (4)	0.500	>0.999	0.679	0.252
Modic change	0.0 (0)	3.6 (2)	0.500	1.8 (1)	3.6 (2)	>0.999	>0.999	>0.999	0.625
Spinal canal stenosis	1.8 (1)	7.1 (4)	0.250	3.6 (2)	3.6 (2)	>0.999	>0.999	0.679	0.243
Kyphosis	30.4 (17)	21.4 (12)	0.267	30.4 (17)	23.2 (13)	0.289	>0.999	>0.999	0.826
Scoliosis	3.6 (2)	3.6 (2)	>0.999	3.6 (2)	1.8 (1)	>0.999	>0.999	>0.999	>0.999

The values are percentages (number of findings per individual) with frequencies (number of findings in group) unless otherwise noted.

Cumulative +Gz stress

The association between spinal injury-induced flight duty limitations and cumulative +Gz exposure in FINAF pilots was studied in 2018.³ The cumulative +Gz stress of the first 5 years of a pilot's career was assessed using a fatigue index. The study population consisted of 23 pilots with spinal injury-induced flight duty limitations and the controls of 50 symptomless pilots. At the time (between 1995 and 2004), the cumulative +Gz load was much higher due to the old fighter training programmes. Thus, our results may be partially explained by the relatively low +Gz exposure during the first 5 years of the pilots' careers. Furthermore, the 5-year follow-up period of the young fighter pilots may be too short to discern a difference between flying and non-flying personnel.

Study evaluation

The strength of present study was the prospective study design and the physically similar groups. In contrast to previous studies, the considerable number of fighter pilots included in this study increased the reliability of our findings. Furthermore, the use of reliable and valid imaging modality (MRI) in this investigation enhances the quality of the study. We classified the MRI findings according to validated methods, which enhances the comparability of our results to other studies.




Figure 1 Boxplot of the intervertebral disc degeneration sum score among different groups.

CONCLUSIONS

In summary, this was a novel prospective study with proper control selection to evaluate the effects of +Gz forces on degenerative changes in the cervical spines of fighter pilots. The result of our study indicates that the effects of +Gz forces on cervical spine degenerative changes were not statistically significant for the most part in comparison with non-flying subjects. The only significant difference between the groups was in the incidence of IVD degeneration of the fighter pilots at C6/C7. It is noteworthy that our fighter pilots were exposed to high +Gz forces for only 3 years during the follow-up period.

The present 5-year follow-up study suggests that significant cervical degeneration is prevalent in young healthy male military personnel. Therefore, further research is needed to ensure that unnecessary neck strain is not applied to fighter pilots or other military personnel.

Acknowledgements We would like to thank Hannu Vähänikkilä, PhD, for advice in statistical analysis.

Contributors The manuscript has been read and approved by all authors. TK has broadly taken part in statistical analysis and drafting and composing the article and also submitted the study. JP has taken big part in drafting and composing the article, especially in the introduction and discussion sections considering previous literature and analysing the data. JK has largely revised and edited the article and taken big part in the interpretation of results. JN took part in defining the study design and reading the MRI's. PO has taken big part in composing the results section and charts and revising the text. TL greatly revised the discussion and conclusions sections and acquired occupational information on pilots. TH was in great role at compiling the study design, in acquisition of data and revising the article.

Funding This study was financially supported by a research grant from the Centre for Military Medicine, Finnish Defence Forces, Finland.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study protocol was approved by the Ethical Committee of the Central Finland Health Care District.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. All data according to Finnish Air Force fighter pilots is considered as classified, and therefore, data used in this study are not available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Sovelius R, Mäntylä M, Heini H, *et al*. Joint helmet-mounted cueing system and neck muscle activity during air combat maneuvering. *Aerosp Med Hum Perform* 2019;90:834–40.
- 2 Coakwell MR, Bloswick DS, Moser R. High-Risk head and neck movements at high G and interventions to reduce associated neck injury. *Aviat Space Environ Med* 2004;75:68–80.
- 3 Honkanen T, Sovelius R, Mäntysaari M, *et al*. +Gz exposure and spinal injury-induced flight duty limitations. *Aerosp Med Hum Perform* 2018;89:552–6.
- 4 Riches A, Spratford W, Witchalls J, *et al*. A systematic review and meta-analysis about the prevalence of neck pain in fast jet pilots. *Aerosp Med Hum Perform* 2019;90:882–90.
- 5 Cheung JPY, Karppinen J, Takatalo J. Cervical degenerative disk disease. In: Shen FH, Samartzis D, Rg F, eds. *Textbook of the cervical spine*. Philadelphia, PA: Elsevier Saunders, 2015: 117–30.
- 6 Shiri R, Frilander H, Sainio M, *et al*. Cervical and lumbar pain and radiological degeneration among fighter pilots: a systematic review and meta-analysis. *Occup Environ Med* 2015;72:145–50.
- 7 Hämäläinen O, Vanharanta H, Kuusela T. Degeneration of cervical intervertebral disks in fighter pilots frequently exposed to high +Gz forces. *Aviat Space Environ Med* 1993;64:692–6.
- 8 Petrén-Mallmin M, Linder J. Mri cervical spine findings in asymptomatic fighter pilots. *Aviat Space Environ Med* 1999;70:1183–8.
- 9 Hendriksen IJ, Holeywijn M. Degenerative changes of the spine of fighter pilots of the Royal Netherlands air force (RNLAf). *Aviat Space Environ Med* 1999;70:1057–63.
- 10 Aydoğ ST, Türbedar E, Demirel AH, *et al*. Cervical and lumbar spinal changes diagnosed in four-view radiographs of 732 military pilots. *Aviat Space Environ Med* 2004;75:154–7.
- 11 Sovelius R, Salonen O, Lamminen A, *et al*. Spinal MRI in fighter pilots and controls: a 13-year longitudinal study. *Aviat Space Environ Med* 2008;79:685–8.
- 12 Tachibana S, Hanada R. Comparative study on F-1 and F-15 pilots spines by MRI in the Japanese air self defence force. In: Burton R, Hämäläinen O, Kuronen P, eds. *Cervical spine injury from repeated exposures to sustained acceleration*. NATO science and technology organization, 1999: 31–2.
- 13 Knudson R, McMillan D, Doucette D, *et al*. A comparative study of G-induced neck injury in pilots of the F/A-18, A-7, and A-4. *Aviat Space Environ Med* 1988;59:758–60.
- 14 McCrary BF, Van Syoc DL. Permanent flying disqualifications of USAF pilots and navigators (1995-1999). *Aviat Space Environ Med* 2002;73:1117–21.
- 15 Honkanen T, Mäntysaari M, Leino T, *et al*. Cross-sectional area of the paraspinal muscles and its association with muscle strength among fighter pilots: a 5-year follow-up. *BMC Musculoskelet Disord* 2019;20:170.
- 16 Miyazaki M, Hong SW, Yoon SH, *et al*. Reliability of a magnetic resonance imaging-based grading system for cervical intervertebral disc degeneration. *J Spinal Disord Tech* 2008;21:288–92.
- 17 Takatalo J, Karppinen J, Niinimäki J, *et al*. Does lumbar disc degeneration on magnetic resonance imaging associate with low back symptom severity in young Finnish adults? *Spine* 2011;36:2180–9.
- 18 Fardon DF, Williams AL, Dohring EJ, *et al*. Lumbar disc nomenclature: version 2.0: recommendations of the combined task forces of the North American spine Society, the American Society of spine radiology and the American Society of Neuroradiology. *Spine J* 2014;14:2525–45.
- 19 Moustarhif M, Bresson B, Koch P, *et al*. Mr imaging of Schmorl's nodes: imaging characteristics and epidemio-clinical relationships. *Diagn Interv Imaging* 2016;97:411–7.
- 20 van der Kraan PM, van den Berg WB. Osteophytes: relevance and biology. *Osteoarthritis Cartilage* 2007;15:237–44.
- 21 Trobisch P, Suess O, Schwab F. Idiopathic scoliosis. *Dtsch Arztebl Int* 2010;107:875–84.
- 22 Harrison DE, Harrison DD, Cailliet R, *et al*. Cobb method or harrison posterior tangent method: which to choose for lateral cervical radiographic analysis. *Spine* 2000;25:2072–8.
- 23 Moll LT, Kindt MW, Stapelfeldt CM, *et al*. Degenerative findings on MRI of the cervical spine: an inter- and intra-rater reliability study. *Chiropr Man Therap* 2018;26:43.
- 24 Matsumoto M, Okada E, Ichihara D, *et al*. Modic changes in the cervical spine: prospective 10-year follow-up study in asymptomatic subjects. *J Bone Joint Surg Br* 2012;94:678–83.
- 25 Oh CH, Yoon SH, Chang HY. Whole spine disc degeneration survey according to the ages and sex using Pfirrmann disc degeneration grades. *Korean J Spine* 2017;14:148–54.